

The radio to X-ray spectrum of GRB 970508 on May 21.0 UT

T.J. Galama¹, R.A.M.J. Wijers², M. Bremer³, P.J. Groot¹, R.G. Strom^{1,4}, C. Kouveliotou^{5,6}, J. van Paradijs^{1,7}

ABSTRACT

We have reconstructed the spectrum of the afterglow of GRB 970508, on May 21.0 UT (12.1 days after the GRB), on the basis of observations spanning the X-ray to radio range. The low-frequency power law index of the spectrum, $\alpha = 0.44 \pm 0.07$ ($F_\nu \propto \nu^\alpha$), is in agreement with the expected value $\alpha = 1/3$ for optically thin synchrotron radiation. The 1.4 GHz emission is self-absorbed. We infer constraints on the break frequencies ν_c and ν_m on May 21.0 UT from a spectral transition from $F_\nu \sim \nu^{-0.6}$ to $F_\nu \sim \nu^{-1.1}$ in the optical passband around 1.4 days. A model of an adiabatically expanding blast wave, emitting synchrotron radiation, in which a significant fraction of the electrons cool rapidly provides a successful and consistent description of the afterglow observations over nine decades in frequency, ranging in time from trigger until several months later.

Subject headings: gamma rays: bursts — gamma rays: individual (GRB 970508)

1. Introduction

GRB970508 was a moderately bright γ -ray burst (Costa et al. 1997; Kouveliotou et al. 1997). It was detected on May 8.904 UT with the Gamma-Ray Burst Monitor (GRBM;

¹Astronomical Institute ‘Anton Pannekoek’, University of Amsterdam, & Center for High Energy Astrophysics, Kruislaan 403, 1098 SJ Amsterdam, The Netherlands

²Institute of Astronomy, Madingley Road, Cambridge, UK

³Institut de Radio Astronomie Millimétrique, 300 rue de la Piscine, F-38406 Saint-Martin d’Hères, France

⁴NFRA, Postbus 2, 7990 AA Dwingeloo, The Netherlands

⁵Universities Space Research Association

⁶NASA/MSFC, Code ES-84, Huntsville AL 35812, USA

⁷Physics Department, University of Alabama in Huntsville, Huntsville AL 35899, USA

Frontera et al. 1991), and with the Wide Field Cameras (WFCs; Jager et al. 1995) on board the Italian-Dutch X-ray observatory BeppoSAX (Piro et al. 1995). Optical observations of the WFC error box (Heise et al. 1997), made on May 9 and 10, revealed a variable object at $RA = 06^h53^m49^s.2$, $Dec = +79^\circ16'19''$ (J2000), which showed an increase by ~ 1 mag in the V band (Bond 1997), and whose spectrum indicated that the distance to the GRB source corresponds to a redshift of at least 0.83 (Metzger et al. 1997). The BeppoSAX Narrow Field Instruments (NFIs) revealed an X-ray transient (Piro et al. 1997a) whose position is consistent with that of the optical variable, and Frail et al. (1997a,b) found the first GRB radio afterglow for GRB 970508; the radio source position coincides with that of the optical source (Bond 1997).

The global properties of the X-ray, optical and radio afterglows of GRBs have been successfully described by relativistic blast wave models (e.g. Wijers, Rees and Mészáros 1997). In this paper we show that the detailed properties of the spectrum of the afterglow of GRB 970508, and its evolution, are well described by a blast wave model in which significant cooling of the electrons plays an important role (Sari, Piran and Narayan 1998). This model predicts the occurrence of an extra break at ν_c in the spectrum, corresponding to the Lorentz factor γ_c above which the electron cooling time is shorter than the expansion time of the blast wave.

2. The X-ray to radio spectrum on May 21.0 UT

In the second week after the GRB 970508 event many observations of its afterglow were made by ground-based and orbiting instruments, which allows us to derive the X-ray to radio afterglow spectrum on May 21.0 UT (12.1 days after the event) with only a few assumptions.

During the first month the 4.86 and 8.46 GHz observations by Frail et al. (1997a) show strong fluctuations attributed to interstellar scintillation (ISS; see e.g. Goodman 1997). At 8.46 GHz the interval of 8 to 14 days after the event corresponds to a flare. It would therefore appear that the ‘true’ underlying 4.86 and 8.46 GHz flux densities are better represented by long-term average values. To discern, apart from an average flux, a possible trend in the 8.46 and 4.86 GHz data of Frail et al. (1997a) we have fitted a power law (in time t , measured in days since the burst) to the data for $t < 30$ days. We find $F_{4.86\text{GHz}} = (101 \pm 17) \times t^{(0.61 \pm 0.06)} \mu\text{Jy}$ ($\chi^2_{\text{red}} = 254/10$) and $F_{8.46\text{GHz}} = (321 \pm 18) \times t^{(0.274 \pm 0.020)} \mu\text{Jy}$ ($\chi^2_{\text{red}} = 521/25$) and we have used these expressions to derive a flux estimate for May 21.0 UT at these frequencies. At lower frequencies the afterglow was not (or barely) detected during this week. The average Westerbork Synthesis Radio Telescope (WSRT) 1.4

GHz flux density (May 9.7-Jul 16.44 UT) is $< 80 \mu\text{Jy}$ (2σ ; Galama et al. 1998a; we note, however, that Frail et al. 1997a report a flux density of $\sim 100 \mu\text{Jy}$).

Pooley and Green (1997) observed GRB 970508 on six consecutive days (May 17-22) with the Ryle telescope at 15 GHz. The source was detected on May 16.52 UT at $1.57 \pm 0.25 \text{ mJy}$, but on none of the other days was an individually significant detection made. However, averaging all observations obtained in the interval May 17-22 (8-13 days after the event), the source was detected at $660 \pm 110 \mu\text{Jy}$. Again, since ISS can modulate the flux densities strongly below $\sim 20 \text{ GHz}$ at this galactic latitude, we assume that the detection on May 16.52 UT was boosted by this effect, and that the source-intrinsic 15 GHz flux on May 21.0 is better represented by the five-day average stated above.

Bremer et al. (1998) detected GRB 970508 with the IRAM Plateau de Bure Interferometer (PdBI) at 86 GHz but not at 232 GHz. No significant variation is observed between the three 86 GHz PdBI detections (May 19.07-May 22.99; center epoch May 21.0 UT) and, combining them we obtain an average $F_{86\text{GHz}} = 1.71 \pm 0.26 \text{ mJy}$. For the 232 GHz upper limit we use the May 21.15 (day 12.2) upper limit of $F_{232\text{GHz}} < 4.22 \text{ mJy}$ (2σ ; Bremer et al. 1998).

In an observation with the Infrared Space Observatory (ISO) on May 21 the source was not detected (Hanlon et al. 1998): preliminary upper limit $F_{12\mu\text{m}} < 120 \mu\text{Jy}$.

We have fitted a power law, $F_K = 120_{-48}^{+75} \times t^{-0.89 \pm 0.27} \mu\text{Jy}$ ($\chi^2_{\text{red}} = 0.23/1$), to the three K band ($2.2 \mu\text{m}$) detections of Chary et al. (1998), and find $F_K = 13_{-8}^{+15} \mu\text{Jy}$ for May 21.0 UT.

For an estimate of the optical flux density we used the fit by Galama et al. (1998b) to the power-law decay of the differential Coussins R (R_c ; $\lambda_c = 6400 \text{ \AA}$) light curve, i.e., $F_{R_c} = (82.7 \pm 2.3) \times t^{-1.141 \pm 0.014} \mu\text{Jy}$; using an absolute calibration uncertainty of 0.1 mag we find $4.8 \pm 0.5 \mu\text{Jy}$.

X-ray fluxes (2-10 keV) of the afterglow, obtained with BeppoSAX, have been reported by Piro et al. (1997b). These observations were made during the burst, and between 6 hours and 6 days afterwards. We have inferred the X-ray flux $F_X = (7.3 \pm 4) \times 10^{-4} \mu\text{Jy}$ for May 21.0 UT by extrapolating the power law fit ($F_X \propto t^{-1.1 \pm 0.1}$) of Piro et al. (1997b).

3. Discussion

3.1. The radio to X-ray spectrum

The radio to X-ray spectrum on May 21.0 UT is shown in Fig. 1. We have divided the spectrum into four parts, corresponding to: the lowest frequencies (I; $< 2.5 \times 10^9$ Hz), the low frequencies (II; $2.5 \times 10^9 - 10^{11}$ Hz), the intermediate frequencies (III; $10^{11} - 10^{14}$ Hz) and the high frequencies (IV; $> 10^{14}$ Hz). Region I has a much steeper spectral slope, $\alpha_{1.4-4.86\text{GHz}} > 1.1$, than region II, $\alpha_{4.86-86\text{GHz}} = 0.44 \pm 0.07$ ($\chi^2_{\text{red}} = 1.4/2$). The 1.4 GHz flux density (region I) is consistent with the expectation for self-absorbed emission ($F_\nu \propto \nu^2$; Katz and Piran 1997), i.e. the self absorption break $\nu_a \sim 2.5$ GHz. The low-frequency region (II) is in agreement with the expected low-frequency tail of synchrotron radiation ($F_\nu \propto \nu^{1/3}$; Rybicki and Lightman 1979). The high frequency X-ray to optical slope ($\alpha = -1.12 \pm 0.07$) is consistent (Galama et al. 1998a; from here on Paper I) with synchrotron radiation from electrons, of which a significant fraction cools rapidly, with a power-law distribution of Lorentz factors, $N(\gamma_e) \propto \gamma_e^{-p}$ (Sari et al. 1998). Optical multi-colour photometry, obtained in the first 5 days after the event, showed that the optical spectrum was well represented by a power law, and approached $\alpha = -1.11 \pm 0.06$ (2.1 to 5.0 days after the GRB; $\chi^2_{\text{red}} = 3.0/4$) after reaching maximum optical light (Galama et al. 1998b; we used $t > 2.1$ days), i.e., consistent with the X-ray to optical slope on May 21.0 UT, suggesting that the slope remained constant over this period. We note that the extrapolation of the X-ray flux is uncertain, but none of the further discussion depends on the validity of the extrapolation, since both spectral breaks we infer lie below the optical. In the spectrum the local optical spectral slope derived from Galama et al. (1998b) is also indicated.

For region III, only upper limits are available. Extrapolating from the adjacent regions and assuming one single spectral break, we find a peak flux of 4.5 mJy at 10^{12} Hz. However, the temporal development of GRB 970508 indicates that a second break frequency exists.

3.2. Evidence for a second spectral break

As noted before by Katz, Piran and Sari (1998), during the first day the optical to X-ray spectral slope of GRB970508 $\alpha \sim -0.5$; which these authors attribute to rapid electron cooling. This is consistent with the UV excess reported by Castro-Tirado et al. (1998), and the reddening of the optical spectrum during the first five days (Galama et al. 1998b). From the data of Galama et al. (1998b) we determine (using $A_V < 0.01$, based on the IRAS $100\mu\text{m}$ cirrus flux, instead of the value $A_V = 0.08$ used by Galama et al. 1998b) $\alpha = -0.54 \pm 0.14$ for t between 0 and 1.5 days, and $\alpha = -1.12 \pm 0.04$ for t between 1.5 and 5 days. This suggests that a spectral break moved through the optical passband between 1.0 and 1.8 days after the burst, separating a range with slope $\alpha \sim -0.5$ from one with

slope $\alpha \sim -1.1$ (see Fig. 2 upper panel). By May 21 we expect this break to have moved downward in frequency to $\sim 10^{14}$ Hz (see Fig. 1 and Sect. 3.4). Additional observational evidence for rapid cooling of a significant fraction of the electrons is given in Paper I.

3.3. Rapid electron cooling

Sari et al. (1998) argued that the most energetic electrons may lose a significant fraction of their energy to radiation. Their model predicts the occurrence of an extra break at ν_c (corresponding to the critical Lorentz factor γ_c above which cooling by synchrotron radiation is significant), in addition to the regular break frequency ν_m (the frequency which corresponds to the lowest-energy injected electrons; Lorentz factor γ_m). The evolution in time of the GRB afterglow is determined by the evolution of the two break frequencies: $\nu_c \propto t^{-1/2}$ and $\nu_m \propto t^{-3/2}$ (Sari et al. 1998). We assume adiabatic evolution of the GRB remnant; see Sect. 3.6. Since ν_m is initially greater than ν_c (Sari et al. 1998), but decays more rapidly, there is a time, t_0 , when the two are equal. For late times $t > t_0$ we have $\nu_m < \nu_c$ and the spectrum varies as $F_\nu \propto \nu^{-(p-1)/2}$ from ν_m up to ν_c ; above ν_c it follows $F_\nu \propto \nu^{-p/2}$ and below ν_m it follows the low-frequency tail, $F_\nu \propto \nu^{1/3}$ (Sari et al. 1998).

3.4. Constraints on the break frequencies ν_c and ν_m for May 21.0 UT

The 86 GHz PdBI observations do not detect the source before 10 days, followed by three detections and subsequently only non detections i.e. at 86 GHz the emission peaked between ~ 10 and ~ 14 days (Bremer et al. 1998). This implies that the peak flux $F_{\nu, \text{max}} \sim 1700 \mu\text{Jy}$ at 86 GHz. The decay after the maximum is quite rapid: for $F_\nu \propto t^\delta$ we find $\delta < -1.1$ using the 3σ upper limit on May 28.41 (Bremer et al. 1998). If this maximum would correspond to the break frequency ν_c passing 86 GHz then we would expect the decay to be much slower, $F_\nu \propto t^{-1/4}$ (Sari et al. 1998). If it reflects the passage of ν_m the subsequent decay would go as $F_\nu \propto t^{3(1-p)/4} = t^{-0.9}$ (Sari et al. 1998). The observed decay rate is in marginal agreement with the passage of the break frequency ν_m , but definitely excludes that ν_c was passing 86 GHz. We therefore identify the maximum at 86 GHz with the passage of ν_m at $t_{\text{m}, 86\text{GHz}} \sim 12$ days, i.e. at the time of the derived spectrum (May 21.0 UT).

There is no evidence for a break at $\nu < 86$ GHz (the 86 GHz detection is consistent with the low-frequency tail; see Fig. 1). This implies that $\nu_c > \nu_m$ and so for $\nu > \nu_m$ the spectrum is predicted to follow $F_\nu \propto \nu^{-(p-1)/2} = \nu^{-0.6}$ (we will use $p = 2.2$, Paper I). Note

also that therefore $t > t_0$ at 12.1 days. The intersection of the high frequency extrapolation with $F_\nu \propto \nu^{-0.6}$ for $\nu > \nu_m$ gives an estimate for ν_c (see Fig. 1); we find $\nu_c = 1.6 \times 10^{14}$ Hz at 12.1 days (May 21.0 UT). Independently, we identify the observed optical spectral transition between 1.0 and 1.8 days from $\alpha = -0.54 \pm 0.14$ to $\alpha = -1.12 \pm 0.04$ (Sect. 3.2 and Fig. 2) with the break frequency ν_c passing the R_c band at t_{c,R_c} . For $t < t_{c,R_c}$ we expect $F_\nu \propto \nu^{-0.6}$, while for $t > t_{c,R_c}$ we expect $F_\nu \propto \nu^{-p/2} = \nu^{-1.1}$ in the optical passband; in excellent agreement with the observed transition. Using $t_{c,R_c} = 1.4$ days ($\nu_c \propto t^{-1/2}$), we find $\nu_c = 1.6 \times 10^{14}$ Hz at $t = 12.1$ days (May 21.0 UT). This is in remarkable agreement with the value derived above from the intersection of $F_\nu \propto \nu^{-0.6}$ for $\nu > \nu_m$ in region III with the high-frequency extrapolation. We therefore identify the transition observed in the optical passband with the passage of the break frequency ν_c at $t_{c,R_c} \sim 1.4$ days. The frequencies ν_a , ν_m and ν_c are shown in Fig. (1); they determine the division of the spectrum into the four regions. We note that the upper limit given for the ISO non detection (Hanlon et al. 1998) is a conservative number and further analysis may provide a stronger constraint on the shape of the spectrum.

3.5. The near-infrared range

At $2.2\mu\text{m}$ (K band) we expect ν_c to pass at $t_{c,K} \sim 16.5$ days ($\nu_c \propto t^{-1/2}$). Before that the K band flux density should decay as $F_\nu \propto t^{-0.9}$ (see above). The decay $F_K \propto t^{-0.89 \pm 0.27}$ (Sect. 3.1), fitted to the data of Chary et al. (1998), is in good agreement with this. The K to R band spectrum should show a gradual transition between 1.4 days (when ν_c passes R_c) and 16.5 days from $F_\nu \propto \nu^{-0.6}$ to $F_\nu \propto \nu^{-1.1}$. Spectral indices α_{K-R_c} , derived from the K band data of Chary et al. (1998) and the optical R_c light curve of Galama et al. (1998b) are in good agreement with this: $\alpha_{K-R_c} = -0.61 \pm 0.12$ ($t = 4.3$), $\alpha_{K-R_c} = -0.69 \pm 0.12$ ($t = 7.3$), and $\alpha_{K-R_c} = -0.91 \pm 0.27$ ($t = 11.3$; see Fig. 2 lower panel). H band ($1.6\mu\text{m}$) HST observations (Pian et al. 1998) detect the OT at $H = 20.6 \pm 0.3$ ($F_H = 6.2 \pm 1.5\mu\text{Jy}$, $t = 24.7$) and comparison with Galama et al. (1998b) gives $\alpha_{H-R_c} = -1.18 \pm 0.29$ (Fig. 2 lower panel). This is consistent with the expectation that ν_c had by then already passed the H band.

3.6. Adiabatic dynamical evolution of the blast wave and the value of t_0

The observed transitions $t_{m,86\text{GHz}} \sim 12$ days and $t_{c,R_c} \sim 1.4$ days imply that $t_0 \sim 0.006$ days (500 s). For $t > t_0$ the dynamical evolution of the remnant is almost certainly adiabatic, while before that time it could possibly be radiative (Sari et al. 1998; but

see Mészáros, Rees & Wijers 1998). This means that one would expect nearly all of the remnant’s observed dynamical evolution to be adiabatic.

However, the maximum flux at 8.46 GHz is $F_{\nu,\text{max}} \sim 700 \mu\text{Jy}$ at $t_{\text{m},8.46\text{GHz}} \sim 55$ days (Frail et al. 1997a); this value of $t_{\text{m},8.46\text{GHz}}$ is as expected from the 86 GHz observations, for $\nu_{\text{m}} \propto t^{-3/2}$. This is less than the peak of $1700 \mu\text{Jy}$ at 86 GHz and would argue for some radiative losses, since in the perfectly adiabatic case the maximum flux should be constant with time, while the peak moves to lower frequencies. Also the rather gradual early evolution of the radio light curves of GRB 970508 and the observed transition from optically thick to thin emission at 1.4 GHz around $t \sim 45$ days suggest the possibility of radiative evolution (Paper I). However, the absence of a break in the smooth power law decay of the optical light curve from 2 to 60 days after the burst (Pedersen et al. 1998; Castro-Tirado et al. 1998; Sokolov et al. 1998; Galama et al. 1998b) shows that there is no important transition in that period. Also the optical spectra and the optical temporal slope indicate adiabatic remnant dynamics with $\nu_{\text{c}} > \nu_{\text{m}}$ during this period (Paper I). Taken together, these results indicate that some additional ingredient is needed to fully explain the radio and mm behaviour; for example, Waxman, Kulkarni and Frail (1998) have argued that the transition from ultrarelativistic to mildly relativistic expansion of the blast wave may explain these anomalies.

4. Conclusion

We have found that on May 21.0 UT the afterglow spectrum of GRB 970508 contains all the characteristic parts of a synchrotron spectrum. We infer that the dynamical evolution of the remnant is adiabatic (in agreement with Waxman et al. 1998), but that a significant fraction of the electrons have synchrotron cooling times shorter than the remnant’s expansion time. In accordance with the theory outlined by Sari et al. (1998) this produces an additional break in the spectrum. The break frequencies ν_{m} and ν_{c} , corresponding to the minimum electron energy and the energy above which electrons cool rapidly, respectively, are equal at a time t_0 which we find is 500 s. This synchrotron spectrum and adiabatic dynamics can explain most of the afterglow behaviour over 9 decades in frequency spanning the X rays to radio waves, from the first optical to the late radio data. Some aspects of the afterglow are not explained by this model: the flux at ν_{m} should be constant, but is seen to decrease from 12 days when it lies at mm wavelengths to 55 days when it lies in the radio; the self-absorption frequency ν_{a} is predicted to be time-independent, but is seen to decrease with time and the early rise of the radio fluxes is slower than expected (see Paper I) and finally the decay after maximum at mm wavelengths is perhaps somewhat faster

than expected (see paper I for a discussion on the radio light curves).

In summary, we find that an adiabatically expanding blast wave emitting synchrotron radiation, with a significant fraction of rapidly cooling electrons, describes most of the afterglow data on GRB 970508 very well.

We would like to thank Dr. Katz for useful discussions. T. Galama is supported through a grant from NFRA under contract 781.76.011. R. Wijers is supported by a Royal Society URF grant. C. Kouveliotou acknowledges support from NASA grant NAG 5-2560.

REFERENCES

- Bond, H. 1997, IAU Circ., 6654
- Bremer, M., Krichbaum, T.P., Galama, T.J., Castro-Tirado, A.J., Frontera, F., van Paradijs, J., Mirabel, I.F., Costa, E. 1998, A&A, 332, L13
- Castro-Tirado, A.J. et al. 1998, Science, 279, 1011
- Chary, R. et al. 1998, accepted for publication in, ApJ
- Costa, E. et al. 1997, IAU Circ., 6649
- Frail, D.A., Kulkarni, S.R., Nicastro, L., Feroci, M., Taylor, G.B. 1997a, Nature, 389, 261
- Frail, D.A. et al. 1997b, IAU Circ., 6662
- Frontera, F. et al. 1991, Adv. Space Res. 11, 281
- Galama, T.J. et al. 1998a, accepted for publication in, ApJ(Paper I)
- Galama, T.J. et al. 1998b, ApJ, 497, 13
- Goodman, J. 1997, New Astronomy, 2, 449
- Hanlon, L. et al. 1998, in preparation (private communication)
- Heise, J. et al. 1997, IAU Circ., 6654
- In 't Zand, J. et al. 1997, IAU Circ., 6569
- Jager, R., Heise, J., In 't Zand, J., Brinkman, A.C. 1995, Adv. Space Res., 13, 315
- Katz, J.I. and Piran, T. 1997, ApJ, 490, 772
- Katz, J.I., Piran, T. and Sari, R. 1998, Phys. Rev. Lett., 80, 1580
- Mészáros, P.I., Rees, M.J., and Wijers, R.A.M.J., 1997, ApJ, in press.

- Metzger, M.R., Djorgovski, S.G., Kulkarni, S.R., Steidel, C.C., Adelberger, K.L., Frail, D.A., Costa, E., Frontera, F. 1997, *Nature*, 387, 878
- Pedersen, H. et al. 1998, *ApJ*, 496, 311
- Pian, E. et al. 1998, *ApJ*, 492, 103
- Piro, L., Scarsi, L. and Butler, R.C. 1995, *Proc. SPIE* 2517, 169
- Piro, L. et al. 1997a, *IAU Circ.*, 6656
- Piro, L. et al. 1997b, *A&A*, 331, L41
- Pooley, G. and Green, D. 1997, *IAU Circ.*, 6670
- Rybicki, G.B., Lightman, A.P. 1979, *Radiative Processes In Astrophysics*, Wiley, New York
- Sari, R., Piran, T. and Narayan, R. 1998, *ApJ*, 497, 17
- Sokolov, V.V., Kopylov, A.I., Zharikov, S.V., Feroci, M., Nicastro, L., Palazzi, E. 1998, accepted for publication, *A&Aastro-ph* 9802341
- Van Paradijs, J. et al. 1997, *Nature*, 386, 686
- Waxman, E., Kulkarni, S.R. & Frail, D.A. 1998, *ApJ*, 497, 288
- Wijers, R.A.M.J., Rees, M.J., and Mészáros, P.I. 1997, *MNRAS*, 288, L51




Fig. 1.— The X-ray to radio spectrum of GRB 970508 on May 21.0 UT (12.1 days after the event). The fit to the low-frequency part, $\alpha_{4.86-86\text{GHz}} = 0.44 \pm 0.07$, is shown as well as the extrapolation from X-ray to optical (solid lines). The local optical slope (2.1–5.0 days after the event) is indicated by the thick solid line. Also indicated is the extrapolation $F_\nu \propto \nu^{-0.6}$ (lines). Indicated are the rough estimates of the break frequencies ν_a , ν_m and ν_c for May 21.0 UT.




Fig. 2.— The spectral slope α as a function of the logarithm of time in days after the burst. The upper panel is from Galama et al. (1998b) and shows the spectral slope transition in the optical passband (slopes determined from U, B, V, R_c and I_c band observations; we used $A_V = 0$, see Sect. 3.2), where data is collected from Galama et al. (1998b; $\star$), Sokolov et al. (1997; $\times$), Djorgovski et al. (1997, $\bullet$), Metzger et al. (1997b, $\circ$) and Castro-Tirado et al (1998; $\square$). The transition from $F_\nu \propto \nu^{-0.6}$ to $F_\nu \propto \nu^{-1.1}$ (dashed lines) occurs at roughly 1.4 days. We expect a similar transition in the spectral slope derived from K- R_c (lower panel) to occur between 1.4 and 16.5 days (see Sect. 3.5). The lower panel shows the spectral indices α_{K-R_c} ($\blacksquare$), derived from the K band data of Chary et al. (1998), and α_{H-R_c} ($\blacktriangle$), derived from the H band data of Pian et al. (1998), with respect to the optical R_c light curve of Galama et al. (1998b).